

Historic, Archive Document

Do not assume content reflects current scientific knowledge, policies, or practices.

1.9422
L8D71B9
Cap.1

DOMESTIC WOOL: MONTHLY AVERAGE PRICE PER POUND, BOSTON MARKET, 1926
GREASE BASIS, OHIO AND SIMILAR

Month	Fine		1/2 Blood		3/8 Blood		1/4 Blood		Low 1/4 blood, strictly		Common and braid	
	Strictly French	Clothing	Strictly French	Clothing	Strictly French	Clothing	Strictly French	Clothing	combing	combing	combing	combing
1926												
Jan.	\$0.530	--	\$0.465	\$0.531	--	\$0.480	\$0.535	\$0.468	\$0.535	\$0.464	\$0.469	\$0.445
Feb.	.520	--	.454	.524	--	.472	.526	.460	.522	--	.461	.440
March	.487	--	.416	.488	--	.435	.490	.441	.487	--	.446	.423
April	.461	--	.405	.460	--	.420	.458	--	.455	--	.442	.395
May	.440	--	.380	.445	--	.398	.438	.380	.436	--	.380	.370
June	.437	--	.365	.439	--	.395	.429	.380	.422	--	.381	.365
46's												
: 64's, 70's, 80's (Fine)		: 58's, 60's (1/2 Blood)		: 56's (3/8 Blood)		: 48's, 50's (1/4 Blood)		: (Low 1/4 Blood)		: 36's, 40's, 44's		
Month	Strictly French	Clothing	Strictly French	Clothing	Strictly French	Clothing	Strictly French	Clothing	Strictly French	Clothing	Strictly French	Clothing
	: combing	: combing	: combing	: combing	: combing	: combing	: combing	: combing	: combing	: combing	: combing	: (Common and braid)
July	\$0.445	--	\$0.366	\$0.440	--	\$0.395	\$0.435	\$0.436	\$0.430	--	\$0.395	\$0.375
Aug.	.445	--	.375	.440	--	.395	.436	.380	.435	--	.395	.375
Sept.	.447	--	.397	.441	--	.395	.440	.383	.436	--	.395	.375
Oct.	.455	--	.405	.452	--	.395	.451	.385	.451	--	.395	.375
Nov.	.455	--	.400	.455	--	.395	.455	.385	.455	--	.395	.375
Dec.	.454	--	.391	.451	--	.390	.451	.385	.449	--	.391	.375
Average	.465	--	.402	.464	--	.414	.462	--	.459	--	.410	.391

Livestock, Meats and Wool Division,
Bureau of Agricultural Economics,
U. S. Department of Agriculture.

LIBRARY
RECEIVED
★ NOV 26 1943
U.S. Department of Agriculture

[illegible]

Abstract

[illegible]

Abstract. The purpose of this study was to determine whether there were differences in the prevalence of periodontitis between patients who had been treated by a general dentist or a periodontist. A total of 60 patients were randomly selected from two dental practices. All patients underwent a clinical examination of their periodontium. The results showed that the prevalence of periodontitis was significantly higher in patients who had been treated by a general dentist than in patients who had been treated by a periodontist. This finding suggests that patients who are treated by a general dentist may have a higher risk of developing periodontitis than those who are treated by a periodontist. Further research is needed to confirm these findings and to identify the factors that contribute to the development of periodontitis.

[illegible]